

## EDITORIAL

- 267** "Obesity-Associated" Breast Cancer in Lean Women: Metabolism and Inflammation as Critical Modifiers of Risk  
Gerald V. Denis and Julie R. Palmer  
*See Cancer Prev Res 10(4):235–43*

## REVIEW

- 270** Sessile Serrated Polyps and Colon Cancer Prevention  
Shahrooz Rashtak, Rafaela Rego, Seth R. Sweetser, and Frank A. Sinicrope

## RESEARCH ARTICLES

- 279** Fatty Acid Synthesis Intermediates Represent Novel Noninvasive Biomarkers of Prostate Cancer Chemoprevention by Phenethyl Isothiocyanate  
Krishna B. Singh and Shivendra V. Singh


- 290** No Decreased Risk of Gastrointestinal Cancers in Users of Metformin in The Netherlands; A Time-Varying Analysis of Metformin Exposure  
Roy G. de Jong, Andrea M. Burden, Sander de Kort, Myrthe P. van Herk-Sukel, Pauline A. Vissers, Paddy K. Janssen, Harm R. Haak, Ad A. Masclee, Frank de Vries, and Maryska L. Janssen-Heijnen

- 298** A Phase II Randomized, Double-blind, Presurgical Trial of Polyphenon E in Bladder Cancer Patients to Evaluate Pharmacodynamics and Bladder Tissue Biomarkers  
Jason R. Gee, Daniel R. Saltzstein, KyungMann Kim, Jill Kolesar, Wei Huang, Thomas C. Havighurst, Barbara W. Wollmer, Jeanne Stublaski, Tracy Downs, Hasan Mukhtar, Margaret G. House, Howard L. Parnes, and Howard H. Bailey

- 308** The Prolyl Isomerase Pin1 Is a Novel Target of 6,7,4'-Trihydroxyisoflavone for Suppressing Esophageal Cancer Growth  
Tae-Gyu Lim, Sung-Young Lee, Zhaoheng Duan, Mee-Hyun Lee, Hanyong Chen, Fangfang Liu, Kangdong Liu, Sung Keun Jung, Dong Joon Kim, Ann M. Bode, Ki Won Lee, and Zigang Dong

## ABOUT THE COVER

Epidemiologically, an inverse correlation has been reported between the intake of soy foods and esophageal cancer, which is the 8th most common cancer worldwide and the 6th most common cause of cancer death with an estimated 400,000 deaths or 4.9% of total cancer deaths. Pin1 is a critical therapeutic and preventive target in esophageal cancer because of its positive regulation of  $\beta$ -catenin and cyclin D1. Indeed, up-regulation of Pin1 levels has been observed clinically and is closely correlated with poor survival of esophageal cancer patients. The cover image reveals the direct binding of 6,7,4'-trihydroxyisoflavone, a major metabolite of daidzein, at both the WW and PPlase domains of Pin1. See article by Lim et al. (beginning on page 308) for more information about this natural inhibitor of the Pin1 protein for suppressing esophageal cancer development.



# Cancer Prevention Research

**10 (5)**

*Cancer Prev Res* 2017;10:267-318.

**Updated version** Access the most recent version of this article at:  
<http://cancerpreventionresearch.aacrjournals.org/content/10/5>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, contact the AACR Publications Department at [permissions@aacr.org](mailto:permissions@aacr.org).