

## EDITORIAL

- 313** Combining Variables for Cancer Risk Estimation: Is the Sum Better than the Parts?  
Christine M. Friedenreich and Anne McTiernan  
*See related article, p. 317*


## RESEARCH ARTICLES

- 317** The Combined Association of Modifiable Risk Factors with Breast Cancer Risk in the Women's Health Initiative  
Rhonda Arthur, Sylvia Wassertheil-Smoller, JoAnn E. Manson, Juhua Luo, Linda Snetselaar, Theresa Hastert, Bette Caan, Lihong Qi, and Thomas Rohan  
*See related editorial, p. 313*
- 327** PAM50 and Risk of Recurrence Scores for Interval Breast Cancers  
Samantha Puvanesarajah, Sarah J. Nyante, Cherie M. Kuzmiak, Mengjie Chen, Chiu-Kit Tse, Xuezheng Sun, Emma H. Allott, Erin L. Kirk, Lisa A. Carey, Charles M. Perou, Andrew F. Olshan, Louise M. Henderson, and Melissa A. Troester

- 337** Inhibition of Glycolysis in Prostate Cancer Chemoprevention by Phenethyl Isothiocyanate  
Krishna B. Singh, Eun-Ryeong Hahm, Lora H. Rigatti, Daniel P. Normolle, Jian-Min Yuan, and Shivendra V. Singh
- 347** Consumption of Sugars, Sugary Foods, and Sugary Beverages in Relation to Adiposity-Related Cancer Risk in the Framingham Offspring Cohort (1991–2013)  
Nour Makarem, Elisa V. Bandera, Yong Lin, Paul F. Jacques, Richard B. Hayes, and Niyati Parekh
- 359** Diagnosing Cervical Neoplasia in Rural Brazil Using a Mobile Van Equipped with *In Vivo* Microscopy: A Cluster-Randomized Community Trial  
Brady Hunt, José Humberto Tavares Guerreiro Fregnani, Richard A. Schwarz, Naitielle Pantano, Suelen Tesoni, Júlio César Possati-Resende, Marcio Antoniazzi, Bruno de Oliveira Fonseca, Graziela de Macêdo Matsushita, Cristovam Scapulatempo-Neto, Ligia Kerr, Philip E. Castle, Kathleen Schmeler, and Rebecca Richards-Kortum

## ABOUT THE COVER

Cervical cancer is a leading cause of death in underserved areas of Brazil, which are often located thousands of kilometers from the nearest facility where diagnostic follow-up and treatment are administered. In the northern part of São Paulo state, Barretos Cancer Hospital has been a pioneer in new mobile strategies for cervical cancer prevention—instead of requiring each patient to travel long distances to a central facility, a mobile vehicular clinic makes routine visits to see patients in their city. While this approach overcomes geographical limitations, new technological innovations are required to administer cervical cancer care in remote locations without compromise in the quality of care. Hunt et al (page 359) report on a cluster randomized trial involving a mobile diagnostic van equipped with a tablet-interfaced *in vivo* microscope. This microscope is a low-cost fiber-optic imaging device that can assess subcellular morphologic features in real time and plays an important role in this mobile strategy. The micrograph featured on the cover of this issue was acquired from a suspected cancer precursor lesion using the *in vivo* microscope and correctly identified by the automated analysis algorithm as requiring treatment.



# Cancer Prevention Research

11 (6)

*Cancer Prev Res* 2018;11:313-370.

**Updated version** Access the most recent version of this article at:  
<http://cancerpreventionresearch.aacrjournals.org/content/11/6>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, use this link <http://cancerpreventionresearch.aacrjournals.org/content/11/6>. Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.