

Perspectives


- Isolating the Effects of Social Interactions on Cancer Biology.** Brian C. Trainor, Colleen Sweeney, and Robert Cardiff843
Perspective on Williams et al., p. 850
- Favoritism in DNA Methylation.** Megan P. Hitchins and Robyn L. Ward847
Perspective on Candiloro and Dobrovic, p. 862

Research Articles

- A Model of Gene-Environment Interaction Reveals Altered Mammary Gland Gene Expression and Increased Tumor Growth following Social Isolation.** J. Bradley Williams, Diana Pang, Bertha Delgado, Masha Kocherginsky, Maria Tretiakova, Thomas Krausz, Deng Pan, Jane He, Martha K. McClintock, and Suzanne D. Conzen850
- Detection of MGMT Promoter Methylation in Normal Individuals Is Strongly Associated with the T Allele of the rs16906252 MGMT Promoter Single Nucleotide Polymorphism.** Ida L.M. Candiloro and Alexander Dobrovic862
- A Pooled Analysis of Continued Prophylactic Efficacy of Quadrivalent Human Papillomavirus (Types 6/11/16/18) Vaccine against High-grade Cervical and External Genital Lesions.** Susanne K. Kjaer, Kristján Sigurdsson, Ole-Erik Iversen, Mauricio Hernandez-Avila, Cosette M. Wheeler, Gonzalo Perez, Darron R. Brown, Laura A. Koutsky, Eng Hseon Tay, Patricia García, Kevin A. Ault, Suzanne M. Garland, Sepp Leodolter, Sven-Eric Olsson, Grace W.K. Tang, Daron G. Ferris, Jorma Paavonen, Matti Lehtinen, Marc Steben, F. Xavier Bosch, Joakim Dillner, Elmar A. Joura, Slawomir Majewski, Nubia Muñoz, Evan R. Myers, Luisa L. Villa, Frank J. Taddeo, Christine Roberts, Amha Tadesse, Janine Bryan, Roger Maansson, Shuang Lu, Scott Vuocolo, Teresa M. Hesley, Alfred Saah, Eliav Barr, and Richard M. Haupt868
- Eleostearic Acid Inhibits Breast Cancer Proliferation by Means of an Oxidation-Dependent Mechanism.** Michael E. Grossmann, Nancy K. Mizuno, Michelle L. Dammen, Todd Schuster, Amitabha Ray, and Margot P. Cleary879
- Urinary Phytoestrogen Excretion and Postmenopausal Breast Cancer Risk: The Multiethnic Cohort Study.** Marc T. Goodman, Yurii B. Shvetsov, Lynne R. Wilkens, Adrian A. Franke, Loic Le Marchand, Kerry K. Kakazu, Abraham M.Y. Nomura, Brian E. Henderson, and Laurence N. Kolonel887
- UVR Exposure Sensitizes Keratinocytes to DNA Adduct Formation.** Sudhir Nair, Vikram D. Kekatpure, Benjamin L. Judson, Arleen B. Rifkind, Richard D. Granstein, Jay O. Boyle, Kotha Subbaramaiah, Joseph B. Guttenplan, and Andrew J. Dannenberg895
- All-Trans Retinoic Acid Suppresses Stat3 Signaling during Skin Carcinogenesis.** Zanobia Syed, Satish B. Cheepala, Jennifer N. Gill, Jennica Stein, Cherie Ann Nathan, John DiGiovanni, Vinita Batra, Patrick Adegboyega, Heather E. Kleiner, and John L. Clifford903
- Obituary**
- Bandaru S. Reddy: In Memoriam (1930–2009).** Chinthalapally V. Rao, Vernon E. Steele, Gary D. Stoner, and Allan H. Conney912

About the Cover

Long-term psychosocial stress is suspected of having adverse effects on breast cancer, but the epidemiologic reports are inconsistent. Williams et al. examined the relationship between the chronic stress of social isolation and mammary-gland carcinogenesis in the FVB-Tg(C3[1]/SV40 T-antigen) mouse model of human breast cancer. After weaning, young female mice were housed for 9.5 weeks in either stressful social isolation or less-stressful grouped housing. The cover image is a stylized depiction of actual testing for the effect of social isolation on mouse behavior in an "open field" testing apparatus consisting of a relatively large area with an object of interest (represented by the red ball) placed several inches from a "home base" container in the lower right corner. Individual mice were placed with some of their own bedding material in the home base and then videotaped for exploratory behavior for five minutes. The movement of a representative isolated mouse (pink track lines) reflects limited exploration, whereas the movement of a representative group-housed mouse (blue track lines) illustrates highly exploratory behavior. Compared with group-housed mice, isolated mice had stress-induced gene expression changes in premalignant mammary-gland tissue (the first such stress-related finding) and developed a significantly larger mammary tumor burden. The actual testing was videotaped in red light because the natural awake cycle of mice is nocturnal, and the tape was analyzed via Ethovision software that allows automated tracking and analysis of an animal's movement. See articles by Williams et al. (beginning on page 850) and Trainor, Sweeney, and Cardiff (beginning on page 843) for more information.



Cancer Prevention Research

2 (10)

Cancer Prev Res 2009;2:843-913.

Updated version	Access the most recent version of this article at: http://cancerpreventionresearch.aacrjournals.org/content/2/10
------------------------	---

E-mail alerts	Sign up to receive free email-alerts related to this article or journal.
Reprints and Subscriptions	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org .
Permissions	To request permission to re-use all or part of this article, use this link http://cancerpreventionresearch.aacrjournals.org/content/2/10 . Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.