

Germline Pathogenic Variants in the Ataxia Telangiectasia Mutated (*ATM*) Gene are Associated with High and Moderate Risks for Multiple Cancers

Michael J. Hall¹, Ryan Bernhisel², Elisha Hughes², Katie Larson², Eric T. Rosenthal², Nanda A. Singh², Johnathan M. Lancaster², and Allison W. Kurian³



ABSTRACT

Pathogenic variants (PVs) in *ATM* are relatively common, but the scope and magnitude of risk remains uncertain. This study aimed to estimate *ATM* PV cancer risks independent of family cancer history. This analysis included patients referred for hereditary cancer testing with a multi-gene panel ($N = 627,742$). Cancer risks for *ATM* PV carriers ($N = 4,607$) were adjusted for family history using multivariable logistic regression and reported as ORs with 95% confidence intervals (CIs). Subanalyses of the c.7271T>G missense PV were conducted. Moderate-to-high risks for pancreatic (OR, 4.21; 95% CI, 3.24–5.47), prostate (OR, 2.58; 95% CI, 1.93–3.44), gastric (OR, 2.97; 95% CI, 1.66–5.31), and invasive ductal breast (OR, 2.03; 95% CI, 1.89–2.19) cancers were estimated for *ATM* PV carriers. Notably, c.7271T>G was associated with higher invasive ductal breast cancer risk (OR, 3.76; 95% CI, 2.76–5.12) than other missense and truncating *ATM* PVs. Low-to-moderate risks were seen for ductal carcinoma *in situ* (OR, 1.80; 95% CI, 1.61–2.02), male breast cancer (OR,

1.72; 95% CI, 1.08–2.75), ovarian cancer (OR, 1.57; 95% CI, 1.35–1.83), colorectal cancer (OR, 1.49; 95% CI, 1.24–1.79), and melanoma (OR, 1.46; 95% CI, 1.18–1.81). *ATM* PVs are associated with multiple cancer risks and, while professional society guidelines support that carriers are eligible for increased breast and pancreatic cancer screening, increased screening for prostate and gastric cancer may also be warranted. c.7271T>G is associated with high risk for breast cancer, with a 3- to 4-fold risk increase that supports consideration of strategies for prevention and/or early detection.

Prevention Relevance: This study estimated risks for multiple cancers associated with *ATM* pathogenic variants independent of family history. These results indicate that some common variants may be associated with higher breast cancer risks than previously appreciated and increased screening for prostate and gastric cancer may be warranted for carriers of *ATM* pathogenic variants.

Introduction

Germline pathogenic variants (PVs) in the ataxia telangiectasia mutated (*ATM*) gene are prevalent (approximately 0.35%) in the population (1, 2). *ATM* PVs are known to be associated with health risks in both an autosomal-dominant and autosomal-recessive fashion. Rare individuals homozygous or compound heterozygous for a germline *ATM* PV develop ataxia telangiectasia (prevalence $\sim 1/40,000$ – $1/100,000$), a childhood-onset disorder typically characterized by global immunodeficiency, cerebellar neurodegeneration, radiation sensitivity, and death in late adolescence (3). Previous estimates of adult-onset

breast cancer risk in women heterozygous for germline *ATM* PVs range from a 2- to 5-fold increased risk compared with women without *ATM* PVs (2, 4–9). It has been suggested that there may be substantially higher risks associated with specific *ATM* variants, such as c.7271T>G (p.Val2424Gly), where the risk of developing breast cancer by age 70 has been estimated at 52%–60% (10–13). Current guidelines from the National Comprehensive Cancer Network (NCCN) advise female *ATM* PV carriers to initiate annual mammographic screening at age 40, with consideration of annual breast MRI, which may be best accomplished at a high-risk breast cancer clinic (14).

Germline *ATM* PVs have also been suggested to increase risks for developing several other adult-onset cancers. Like the increased risk of breast cancer, these are generally regarded as low to moderately increased lifetime cancer risks. Multiple studies have documented an association of germline *ATM* PVs with pancreatic cancer, although risk estimates vary (15, 16). There is limited evidence for additional malignancies including gastro-esophageal cancer, (17, 18) colorectal cancer (19, 20), ovarian cancer (7, 21), melanoma (22, 23), prostate cancer (22, 24), thyroid cancer (22), gastric cancer (17), and head and neck cancers (22). Estimates of the varied cancer risks associated with *ATM* PVs have been limited by studies with small participant sample sizes leading to imprecise estimates of

¹Fox Chase Cancer Center, Philadelphia, Pennsylvania. ²Myriad Genetics, Inc., Salt Lake City, Utah. ³Stanford University, Stanford, California.

Note: Supplementary data for this article are available at Cancer Prevention Research Online (<http://cancerprevres.aacrjournals.org/>).

Corresponding Author: Michael J. Hall, Cancer Prevention and Control Program, Fox Chase Cancer Center, Philadelphia, PA 19111. Phone: 215-728-2861; E-mail: michael.hall@fccc.edu

Cancer Prev Res 2021;14:433–40

doi: 10.1158/1940-6207.CAPR-20-0448

©2021 American Association for Cancer Research.

Hall et al.

cancer risks, sequencing of limited loci, and ascertainment biases in selection of participants.

Improved precision of estimates of cancer risks associated with *ATM* PVs is critical because *ATM* PVs are common in individuals of varied ancestry undergoing genetic testing for hereditary cancer risk (25, 26), and even modest risk increases may warrant modified recommendations for cancer screening and prevention. We used multi-gene sequencing results and personal and family cancer history data from a large sample of individuals undergoing hereditary cancer risk evaluation to obtain ORs and penetrance estimates, adjusted for family history, for many cancer types. The differential impact of mutation type (nonsense/frameshift versus missense PVs), and the previously described c.7271T>G missense variant were specifically examined.

Penetrance estimates adjusted for family history represent the magnitude of genetic risk that is independent of family history. Adjusted penetrance may be estimated in an unbiased manner from clinical populations where factors related to ascertainment are well captured (7, 27, 28). These adjusted penetrance estimates are applicable to women with and without family history of cancer and may inform personalized assessments that combine genetic risks with environmental and lifestyle risk factors.

Methods

Cohort and hereditary cancer testing

The study included individuals who underwent testing with a multi-gene hereditary cancer panel including *ATM* between September 2013 and July 2019. All tested individuals were ascertained by healthcare providers for suspicion of hereditary cancer risk based on personal and/or family histories of cancer meeting professional society criteria for genetic testing. During this time, the number of genes on the panel expanded from 25 (2013–2016) to 29 (2016–2019) to 35 (2019). Individuals were eligible for analysis if they carried a single (heterozygous) PV in *ATM* or if they had no PV in any gene sequenced nor any uncertain variant (VUS) in the *ATM* gene. Variant classification was performed by the testing laboratory based on American College of Medical Genetics guidelines at a minimum, as described previously (29). Variants with a laboratory classification of deleterious or suspected deleterious were considered to be PVs for this analysis.

Genetic testing data were correlated with demographic, personal, and family cancer history data from a test requisition form that accompanied each test kit. Individuals with missing data on gender or age of testing were excluded. Finally, patients from states that restrict the use of deidentified genetic data after completion of testing were excluded from analysis.

Statistical methods

All analyses were performed using R software (R Foundation for Statistical Computing; version 3.6.1). Multivariable logistic regression modeling was used to quantify *ATM* cancer risks for 11 different cancers in terms of ORs adjusted for age, ancestry,

and personal and family cancer history. These ORs are a measure of cancer risks associated with *ATM* PVs and can be interpreted as relative risks independent of family history (e.g., an OR of 2 indicates that risk is doubled for a PV carrier versus a noncarrier with an identical family history of cancer). A separate analysis was performed for each *ATM* PV subgroup [all *ATM*, truncating (nonsense and frameshift), missense, and the c.7271T>G PV] using PV status as the dependent variable. Independent variables included age, sex, ancestry, and personal and family cancer histories associated with hereditary breast and ovarian cancer, and Lynch and adenomatous polyposis cancer syndromes. For history of breast cancer, categories included ductal invasive (coded as either triple-negative or hormone-positive), lobular invasive, ductal carcinoma *in situ* (DCIS), and male breast. Triple-negative breast cancer is a subset of ductal invasive for this cohort. All other categories are exclusive. Personal cancer variables were coded as binary (ever or never affected). Familial cancers were coded as numeric counts of diagnoses, weighted by degree of relatedness.

Cumulative invasive breast cancer risks were calculated according to the product-limit method (30) without adjustment for competing mortality. These calculations have been described in detail previously (7, 31). Briefly, age-specific incidence rates were estimated as products of ORs and general population incidences; ORs for PVs were calculated as specified above and were combined with age-specific SEER incidence rates from 2009 to 2013 in 5-year intervals to estimate absolute cancer risks (7, 32, 33).

Confidence intervals (CIs) and *P* values were calculated using Wald statistics. All *P* values are reported as two-sided. The study was reviewed by the Fox Chase Cancer Center Institutional Review Board and was determined to be Human Subject Exempt and therefore written informed consent was not required.

Results

We identified 627,742 eligible patients (Table 1), including 4,607 (0.7%) individuals with an *ATM* PV and 623,135 (99.3%) individuals with no *ATM* PV, *ATM* VUS, or PV in another hereditary cancer risk gene. The majority of *ATM* PVs identified were truncating (nonsense and frameshift; 58.3%) and the remaining were missense (7.1%) or other (29.7%). The c.7271T>G variant was the most frequent individual PV observed (5.0%, 229/4607). All observed PV types and their locations on *ATM* are shown in Fig. 1.

The study cohort was predominantly female (96.5%) and over half of individuals reported White (non-Hispanic) ancestry (57.5%; Table 1). Median age of testing was 58 years (range 6, ≥90) for the full cohort. More than one-third (38.2%) of individuals reported a personal history of at least one cancer, and patients with an *ATM* PV were proportionally more likely to report a personal history of cancer than those without an *ATM* PV. Ductal invasive breast cancer was the most common cancer reported. Similarly, ductal invasive breast cancer was by

Table 1. Demographics and personal cancer history by *ATM* PV carrier status.

Characteristic	<i>ATM</i> PV (N = 4,607)	No <i>ATM</i> PV ^a (N = 623,135)	Total (N = 627,742)
Age at testing - median (IQR)	49 (39, 59)	48 (38, 58)	48 (38, 58)
Sex			
Female	4,378 (95.0)	601,246 (96.5)	605,624 (96.5)
Male	229 (5.0)	21,889 (3.5)	22,118 (3.5)
Race/Ethnicity			
White/Caucasian	2,889 (62.9)	358,097 (57.5)	360,986 (57.5)
Black/African	289 (6.3)	51,324 (8.2)	51,613 (8.2)
Latino/Hispanic	288 (6.3)	46,713 (7.5)	47,001 (7.5)
Asian	100 (2.2)	14,224 (2.3)	14,324 (2.3)
Ashkenazi Jewish	34 (0.7)	10,302 (1.7)	10,336 (1.6)
Native American	31 (0.7)	5,141 (0.8)	5,172 (0.8)
Middle Eastern	21 (0.5)	3,428 (0.6)	3,449 (0.5)
Multiple	270 (5.9)	41,021 (6.6)	41,291 (6.6)
Other/None specified	675 (14.7)	92,885 (14.9)	93,560 (14.9)
Personal cancer history			
Total cancers			
0 cancer	2,243 (48.7)	385,932 (61.9)	388,175 (61.8)
1 cancer	1,928 (41.8)	203,668 (32.7)	205,596 (32.8)
2+ cancers	436 (9.5)	33,535 (5.4)	33,971 (5.4)
Any Cancer	2,364 (51.3)	237,203 (38.1)	239,567 (38.2)
Breast - Ductal Invasive ^b	1,461 (33.4)	132,465 (22.0)	133,926 (22.1)
Breast - Triple-negative ^b	52 (1.2)	19,711 (3.3)	19,763 (3.3)
Breast - Lobular Invasive ^b	68 (1.6)	10,633 (1.8)	10,701 (1.8)
Breast - DCIS ^b	351 (8.0)	28,906 (4.8)	29,257 (4.8)
Breast - Male ^c	20 (0.7)	1,909 (0.7)	1,929 (0.7)
Ovarian cancer ^b	192 (4.4)	24,616 (4.1)	24,808 (4.1)
Colorectal cancer	135 (2.9)	14,527 (2.3)	14,662 (2.3)
Endometrial cancer ^b	85 (1.9)	11,877 (2.0)	11,962 (2.0)
Melanoma	88 (1.9)	7,426 (1.2)	7,514 (1.2)
Prostate cancer ^c	75 (32.8)	4,436 (20.3)	4,511 (20.4)
Pancreatic cancer	64 (1.4)	2,085 (0.3)	2,149 (0.3)
Gastric cancer	12 (0.3)	518 (0.1)	530 (0.1)

^aPatients with no PVs in any gene tested, including *ATM*, and no VUS in *ATM*.

^bFemale only.

^cMale only.

Triple-negative breast cancer is a subset of ductal invasive for this cohort. All other categories are exclusive.

far the most common family history reported, followed by ovarian and colorectal cancers (Supplementary Table S1).

Because previous research has identified variability in breast cancer risk by mutation type, with several studies reporting high breast cancer risks associated with the missense *c.7271T>G* PV in *ATM*, we examined rates of personal and family history of breast cancer reported by individuals with a truncating (nonsense and frameshift) PVs, missense PVs, and the *c.7271T>G* missense PV (Supplementary Table S2). Among carriers of *ATM* PVs, 38.7% (120/310) of missense mutation carriers, 45.3% (101/223) of *c.7271T>G* PV carriers, and 32.8% (837/2548) of truncating (frameshift and nonsense) PV carriers had a personal history of invasive breast cancer. No differences in age of cancer diagnosis were detected by mutation.

Multivariable models were used to estimate the family history-adjusted risk of developing cancer in the presence of any *ATM* PV. Results for any *ATM* PV are reported in Table 2 and show a 2-fold increased risk (OR, 2.03; 95% CI, 1.89–2.19)

of ductal invasive breast cancer for women, a 1.80-fold increased risk of DCIS (95% CI, 1.61–2.02), and a 1.72-fold increased risk of male breast cancer (95% CI, 1.08–2.75). Increased risks of several other cancers were also seen in *ATM* PV carriers compared with noncarriers. The largest increase in risk was for pancreatic cancer (OR, 4.21; 95% CI, 3.24–5.47), but gastric cancer (OR, 2.97; 95% CI, 1.66–5.31) and prostate cancer (OR, 2.58; 95% CI, 1.93–3.44) also showed marked elevations in risk. The risks of ovarian cancer (OR, 1.57; 95% CI, 1.35–1.83), colorectal cancer (OR, 1.49; 95% CI, 1.24–1.79), and melanoma (OR, 1.46; 95% CI, 1.18–1.81) were overall low-to-moderately increased among *ATM* PV carriers.

Family history-adjusted risk of breast cancer was also examined by mutation type and specifically in carriers of the *c.7271T>G* missense PV (Supplementary Table S3). The number of *c.7271T>G* carriers was insufficient to generate variant-specific risk estimates for cancers other than breast. Missense *ATM* PVs (OR, 2.52; 95% CI, 1.94–3.28) and truncating (nonsense and frameshift) *ATM* PVs (OR, 1.95; 95% CI,

Hall et al.




Figure 1.

Map of *ATM* and all variants discovered in this study. This map shows the types and frequency of *ATM* variants discovered within this patient cohort. The circle size/line length reflects the number of times a variant was observed at that location. The exons are numbered, with in-frame exons being dark blue and out-of-frame exons being gray. The untranslated region is reflected in light blue. The functional regions of the gene are also shown.

1.77–2.14) were associated with similar risks of ductal invasive breast cancer. Risk estimates were significantly higher for c.7271T>G carriers (OR, 3.76; 95% CI, 2.76–5.21) compared with all other *ATM* PVs. Nonsense/frameshift PVs were also associated with an increased risk of DCIS (OR, 2.05; 95% CI, 1.78–2.36). The c.7271T>G PV was associated with an increased risk of DCIS (OR, 1.70; 95% CI, 1.03–2.81) and male breast cancer (OR, 8.31; 95% CI, 1.46–47.27). Cumulative risk

of developing invasive breast cancer compared with population risk was modeled over an 80-year lifespan separately for each mutation type (Fig. 2).

Discussion


These results provide comprehensive estimates of the multiple cancer risks attributable to PVs in *ATM*, including close to 2-fold elevated risks of invasive breast cancer and DCIS in female PV carriers. In addition, moderate to high risks of pancreatic cancer (greater than 4-fold increase), gastric cancer (3-fold increase), and prostate cancer (2- to 3-fold increase), and modestly yet significantly elevated risk of several other common tumors including colorectal cancer (1.5-fold increase) were observed. Of the cancers evaluated, only endometrial cancer did not show evidence of significant association with *ATM* PVs. Furthermore, we demonstrate through analyses of the common c.7271T>G PV that there is site-specific variability in cancer risks with *ATM* PVs, constituting a genotype-phenotype association. While not all previously reported risks were confirmed by these analyses due to small numbers of reported cancers (e.g., thyroid cancer, head and neck cancers), it remains possible that such associations might be detectable in a larger, more diverse sample. Our estimates build on previous studies suggesting multiple risks of cancer for *ATM* PV carriers, (17, 22) and offer new precision to the magnitude of those risks after controlling for confounding factors.

Table 2. ORs for cancer risk for all *ATM* PVs.

Cancer type	All <i>ATM</i> PVs	
	OR (95% CI)	P
Breast - Ductal Invasive	2.03 (1.89–2.19)	<0.0001
Breast - DCIS	1.80 (1.61–2.02)	<0.0001
Breast - Lobular Invasive	0.94 (0.74–1.20)	0.6229
Breast - Male	1.72 (1.08–2.75)	0.0233
Ovarian	1.57 (1.35–1.83)	<0.0001
Colorectal	1.49 (1.24–1.79)	<0.0001
Endometrial	1.10 (0.88–1.36)	0.4079
Melanoma	1.46 (1.18–1.81)	0.0006
Prostate	2.58 (1.93–3.44)	<0.0001
Pancreatic	4.21 (3.24–5.47)	<0.0001
Gastric	2.97 (1.66–5.31)	0.0002

Note: ORs are adjusted for personal (coded as binary, affected/unaffected, variables) and family (numeric counts weighted by degree of relation) cancer histories of breast (invasive, ductal, DCIS, and male), ovarian, colorectal, melanoma, gastric, pancreatic, prostate, endometrial, and colon (polyps), as well as age, sex, and ancestry.

Figure 2.
Invasive breast cancer cumulative risk by variant type. Groups except for All are exclusive.



The importance of precise and comprehensive estimates of cancer risks associated with *ATM* PVs is supported by the high prevalence of *ATM* PVs in the general population (1, 2), the recognition of variability in cancer risk by PV type and position (10–13), and the variety of common adult tumors like breast cancer, colorectal cancer, and prostate cancer whose risks may be moderately to highly increased by a germline *ATM* PV. It has been 30 years since Swift first published data detailing an increased risk of cancer in Ataxia-Telangiectasia families, (2) and understanding of the genetic risks underlying common cancers has advanced enormously. As genetic testing uptake increases, the identification of germline *ATM* PVs may contribute to reducing cancer incidence across a number of disease sites. Moreover, the development of PARP inhibitors for tumors with homologous recombination deficiency due to disrupted function of *BRCA1/2* and other Fanconi Anemia pathway-associated genes suggests that germline and tumor testing for *ATM* PVs may ultimately guide treatment selection (34, 35).

Previous studies have demonstrated variability in breast cancer risk depending on the type of *ATM* PV (truncating versus missense; ref. 6). The *c.7271T>G* missense variant constituted approximately 5% of all *ATM* PVs identified in this study. Previous research has strongly suggested that this variant increases cancer risk through a dominant-negative effect causing higher cancer risks in *c.7271T>G* carriers compared with other *ATM* PVs (10, 12, 13, 36–38). Our estimate of family history-adjusted ductal invasive breast cancer risk associated with the *c.7271T>G* PV (OR, 3.76; 95% CI, 2.76–5.12) is similar to that estimated for *BRCA2* PVs using the same methodology (7). The family history-adjusted estimates presented here should be slightly lower than previously published

unadjusted estimates, because the risk attributable to family history and *ATM* PV status is not double-counted. However, our estimates fall within the ranges reported from previous studies, all of which had wide CIs. These include estimates from Goldgar and colleagues (OR, 8.0; 95% CI, 2.3–27.4; ref. 12), Southey and colleagues (OR, 11.0; 95% CI, 1.4–85.7; ref. 11), Bernstein and colleagues (8.6-fold increased risk; 95% CI, 3.9–18.9-fold; ref. 13), and Chenevix-Trench and colleagues (13.7-fold increased risk; 95% CI, 5.1–36.6-fold; ref. 10). Compared with these other studies, our study offers the advantage of a methodology that has been shown to produce unbiased results, and 10-fold more individuals genotyped and found to carry *c.7271T>G* noncarriers. *ATM* is a large gene comprised of 62 coding exons and 3,056 amino acids (Fig. 1), and plays a variety of roles in cellular processes including regulation of DNA damage response, DNA repair through multiple pathways, genotoxic stress response, and others (3). Therefore, even rarer high-risk PVs like *c.7271T>G* may be scattered throughout this complex gene, which have, to date, eluded discovery.

These results demonstrate that *ATM* PV carriers have increased risk of pancreatic cancer (OR, 4.21), prostate cancer (OR, 2.58), and gastric cancer (OR, 2.97). *ATM* PV carriers were not included in the initial high-risk guidelines for pancreatic cancer published by the International Cancer of the Pancreas Screening (CAPS) Consortium in 2013 (39), but since that time evidence has emerged demonstrating that germline *ATM* PVs are the most common, or second most common after *BRCA2* PVs, findings in pancreatic cancer patients (15, 40). A recent study estimated a 5.7 OR for pancreatic cancer in *ATM* PV carriers, very close to what was calculated in the same study for *BRCA2* (OR, 6.2; ref. 15). Although *ATM* PV carriers are

Hall et al.

included in the CAPS5 pancreatic cancer screening study (NCT02000089), they are treated as lower risk than carriers of PVs in *BRCA2* and *PALB2*. Future study design should consider the updated relative risk estimates for *ATM* from this and other studies.

The high prevalence of prostate cancer in the United States population, as well as the challenge of differentiating prostate cancers associated with high morbidity and mortality from more indolent tumors, highlights the importance of identifying molecular and genetic factors associated with risk, treatment, and outcomes. The prostate cancer history data in this analysis did not include Gleason score, so we could not determine whether *ATM* PVs are associated with more aggressive disease. Germline *ATM* PVs have previously been associated with more aggressive prostate cancers and shown to be susceptible to PARP inhibitor therapy (35, 41). Finally, our estimates of gastric cancer risk are consistent with those of Helgason and colleagues who estimated 4- to 5-fold increased risks of gastric cancer (17), often early-onset, in carriers of germline loss-of-function *ATM* PVs. Though gastric cancer is rare in the United States (~28,000 new cases annually), morbidity is exceedingly high (5-year survival ~30%; ref. 32), highlighting the value of risk stratification through predictive genetic testing to promote risk factor mitigation (e.g., treatment of *H. pylori* infection, smoking cessation) and determine potential eligibility for gastric cancer screening, as performed in individuals with Lynch syndrome (42). Notably, no differences in age at diagnosis were observed for prostate and gastric cancers based on whether an *ATM* PV was present or not. Further exploration of how *ATM* PV status may affect current screening guidelines is warranted for both of these tumors.

A recently published study examining data from a large sample of patients with diverse personal and cancer histories tested by multigene panel testing through a single laboratory reported ORs for *ATM* PV carriers for breast cancer (OR, 2.91; 95% CI, 2.48–3.42), ovarian cancer (OR, 1.91; 95% CI, 1.37–2.65), pancreatic cancer (OR, 5.88; 95% CI, 3.64–9.4), colorectal cancer (OR, 2.6; 95% CI, 1.53–4.3), endometrial cancer (OR, 1.44; 95% CI, 0.62–3.2), and melanoma (OR, 2.69; 95% CI, 1.32–5.23; ref. 43). Methodologies were notably different from those employed in our study, with the authors comparing *ATM* PVs in Caucasian patients with breast cancer referred for hereditary cancer testing against PV frequencies observed in non-Finnish European controls from the gnomAD database. This methodology may significantly overestimate risk, because clinically ascertained cases are likely to be enriched for mutations over general population cases (44). Despite this methodologic difference, risk estimates are similar to those we report here.

Our study has several important limitations. Personal and family cancer history data were collected from a standard test requisition form; thus cancer pathology and age of diagnosis have not been verified. However, we have previously shown through sensitivity analyses that over- or under-reporting of family history would have modest or no impact on the risk

estimates calculated using multivariable logistic regression (7). In addition, despite the large sample size examined here, the relative rarity of missense *ATM* PVs and the c.7271T>G PV limited statistical power to estimate risks for cancer types beyond breast cancer. Similarly, small samples limited statistical power to estimate cancer risks precisely in rarer tumor types, such as triple-negative breast cancer and male breast cancer. In addition, our sample is notably female predominant (97%) reflective of the historically strong association of genetic testing for hereditary cancer risk with female breast cancer despite the broadening of testing indications in recent years to include prostate cancer and male-dominant cancers such as pancreatic cancer. In addition, we may not have been able to detect situations where the risks for certain cancers, for example, pancreatic, differ between men and women. Nonetheless, it should be noted that our proportionally smaller male population includes over 22,000 male patients and 229 male *ATM* PV carriers, as well as family history data in both male and female relatives among study participants. In conclusion, through analyses including multivariable modeling in a very large sample of individuals undergoing multi-gene panel testing for hereditary cancer risk, we demonstrate that PVs in *ATM* are associated with moderate- to high-risks for multiple adult-onset cancers, independent of family history. High risks of prostate, gastric, and notably pancreatic cancer may warrant screening in *ATM* PV carriers. Furthermore, these results confirm that the c.7271T>G missense PV is associated with high risk for ductal invasive ductal breast cancer, with a three- to 4-fold risk increase that is similar to that of *BRCA2* PVs and supports consideration of similar strategies for prevention and/or early detection.

Authors' Disclosures

M.J. Hall reports several collaborative research studies with Myriad. These studies are purely collaborative in nature and have not involved any compensation or funding to M.J. Hall from Myriad. R. Bernhisel reported personal fees from Myriad Genetics, Inc. during the conduct of the study. E. Hughes reported personal fees from Myriad Genetics, Inc. during the conduct of the study. E.T. Rosenthal reported personal fees from Myriad Genetics, Inc. during the conduct of the study. N.A. Singh reported personal fees from Myriad Genetics, Inc. during the conduct of the study. J.M. Lancaster is currently an employee of Regeneron Pharmaceuticals but received personal fees from Myriad Genetics, Inc. during the conduct of the study. A.W. Kurian reported grants from Myriad Genetics outside the submitted work. No other disclosures were reported.

Authors' Contributions

M.J. Hall: Conceptualization, visualization, methodology, writing—original draft, writing—review and editing. **R. Bernhisel:** Resources, formal analysis, validation, visualization, methodology, writing—review and editing. **E. Hughes:** Resources, formal analysis, validation, visualization, methodology, writing—review and editing. **K. Larson:** Resources, formal analysis, validation, methodology, writing—review and editing. **E.T. Rosenthal:** Conceptualization, resources, funding acquisition, visualization, methodology, writing—review and editing. **N.A. Singh:** Conceptualization, resources, funding acquisition, visualization, writing—review and editing. **J.M. Lancaster:** Conceptualization, supervision,

Germline ATM PVs Are Associated with Multiple Cancer Risks

funding acquisition, writing–review and editing. **A.W. Kurian:** Conceptualization, visualization, methodology, writing–original draft, writing–review and editing.

Acknowledgments

The authors would like to acknowledge Stephanie Meek and Gabby Iorg for their assistance in manuscript preparation.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received September 9, 2020; revised November 20, 2020; accepted January 21, 2021; published first January 28, 2021.

References

- Swift M, Morrell D, Cromartie E, Chamberlin AR, Skolnick MH, Bishop DT. The incidence and gene frequency of ataxia-telangiectasia in the United States. *Am J Hum Genet* 1986;39:573–83.
- Swift M, Reitnauer PJ, Morrell D, Chase CL. Breast and other cancers in families with Ataxia-Telangiectasia. *N Engl J Med* 1987; 316:1289–94.
- Rothblum-Oviatt C, Wright J, Lefton-Greif MA, McGrath-Morrow SA, Crawford TO, Lederman HM. Ataxia telangiectasia: a review. *Orphanet J Rare Dis* 2016;11:159.
- Swift M, Morrell D, Massey RB, Chase CL. Incidence of cancer in 161 families affected by Ataxia-Telangiectasia. *N Engl J Med* 1991;325: 1831–6.
- Renwick A, Thompson D, Seal S, Kelly P, Chagtai T, Ahmed M, et al. ATM mutations that cause ataxia-telangiectasia are breast cancer susceptibility alleles. *Nat Genet* 2006;38:873–5.
- Tavtigian SV, Oefner PJ, Babikyan D, Hartmann A, Healey S, Le Calvez-Kelm F, et al. Rare, evolutionarily unlikely missense substitutions in ATM confer increased risk of breast cancer. *Am J Hum Genet* 2009;85:427–46.
- Kurian AW, Hughes E, Handorf EA, Gutin A, Allen B, Hartman A-R, et al. Breast and ovarian cancer penetrance estimates derived from germline multiple-gene sequencing results in women. *JCO Precis Oncol* 2017;1:1–12.
- Hauke J, Horvath J, Groß E, Gehrig A, Honisch E, Hackmann K, et al. Gene panel testing of 5589 BRCA1/2-negative index patients with breast cancer in a routine diagnostic setting: results of the German Consortium for Hereditary Breast and Ovarian Cancer. *Cancer Med* 2018;7:1349–58.
- Lu H-M, Li S, Black MH, Lee S, Hoiness R, Wu S, et al. Association of breast and ovarian cancers with predisposition genes identified by large-scale sequencing. *JAMA Oncol* 2019;5:51–7.
- Chenevix-Trench G, Spurdle AB, Gatei M, Kelly H, Marsh A, Chen X, et al. Dominant negative ATM mutations in breast cancer families. *J Natl Cancer Inst* 2002;94:205–15.
- Southey MC, Goldgar DE, Winqvist R, Pylkäs K, Couch F, Tischkowitz M, et al. PALB2, CHEK2 and ATM rare variants and cancer risk: data from COGS. *J Med Genet* 2016;53:800–11.
- Goldgar DE, Healey S, Dowty JG, Da Silva L, Chen X, Spurdle AB, et al. Rare variants in the ATM gene and risk of breast cancer. *Breast Cancer Res* 2011;13:R73.
- Bernstein JL, Teraoka S, Southey MC, Jenkins MA, Andrulis IL, Knight JA, et al. Population-based estimates of breast cancer risks associated with ATM gene variants c.7271T>G and c.1066–6T>G (IVS10–6T>G) from the Breast Cancer Family Registry. *Hum Mutat* 2006;27:1122–8.
- Daly MB, Pilarski R, Berry M, Buys SS, Friedman S, Garber JE, et al. NCCN Clinical practice guidelines in oncology, genetic/familial high-risk assessment: breast and ovarian (Version 3.2019). In: NCCN Clinical Practice Guidelines in Oncology; 2019. Available from: https://www.nccn.org/professionals/physician_gls/pdf/genet_ics_bop.pdf.
- Hu C, Hart SN, Polley EC, Gnanaolivu R, Shimelis H, Lee KY, et al. Association between inherited germline mutations in cancer predisposition genes and risk of pancreatic cancer. *JAMA* 2018;319:2401–9.
- Shindo K, Yu J, Suenaga M, Fesharakizadeh S, Cho C, Macgregor-Das A, et al. Deleterious germline mutations in patients with apparently sporadic pancreatic adenocarcinoma. *J Clin Oncol* 2017;35:3382–90.
- Helgason H, Rafnar T, Olafsdottir HS, Jonasson JG, Sigurdsson A, Stacey SN, et al. Loss-of-function variants in ATM confer risk of gastric cancer. *Nat Genet* 2015;47:906–10.
- Huang D-S, Tao H-Q, He X-J, Long M, Yu S, Xia Y-J, et al. Prevalence of deleterious ATM germline mutations in gastric cancer patients. *Oncotarget* 2015;6:40953–8.
- Pearlman R, Frankel WL, Swanson B, Zhao W, Yilmaz A, Miller K, et al. Prevalence and spectrum of germline cancer susceptibility gene mutations among patients with early-onset colorectal cancer. *JAMA Oncol* 2017;3:464–71.
- Yurgelun MB, Allen B, Kaldete RR, Bowles KR, Judkins T, Kaushik P, et al. Identification of a variety of mutations in cancer predisposition genes in patients with suspected lynch syndrome. *Gastroenterology* 2015;149:604–13.
- Kurian AW, Ward KC, Howlader N, Deapen D, Hamilton AS, Mariotto A, et al. Genetic testing and results in a population-based cohort of breast cancer patients and ovarian cancer patients. *J Clin Oncol* 2019;37:1305–15.
- Dombernowsky SL, Weischer M, Allin KH, Bojesen SE, Tybjjrg-Hansen A, Nordestgaard BG. Risk of cancer by ATM missense mutations in the general population. *J Clin Oncol* 2008;26:3057–62.
- Goldstein AM, Xiao Y, Sampson J, Zhu B, Rotunno M, Bennett H, et al. Rare germline variants in known melanoma susceptibility genes in familial melanoma. *Hum Mol Genet* 2017;26:4886–95.
- Pritchard CC, Mateo J, Walsh MF, De Sarkar N, Abida W, Beltran H, et al. Inherited DNA-repair gene mutations in men with metastatic prostate cancer. *N Engl J Med* 2016;375:443–53.
- Rosenthal ET, Bernhisel R, Brown K, Kidd J, Manley S. Clinical testing with a panel of 25 genes associated with increased cancer risk results in a significant increase in clinically significant findings across a broad range of cancer histories. *Cancer Genet* 2017;218–219:58–68.
- Couch FJ, Shimelis H, Hu C, Hart SN, Polley EC, Na J, et al. Associations between cancer predisposition testing panel genes and breast cancer. *JAMA Oncol* 2017;3:1190–6.
- Rothman KJ, Greenland S, Lash T. *Modern epidemiology*. Philadelphia, PA: Lippincott Williams & Wilkins; 2008:95–7.
- Rajamani S. Eliminating bias in cancer risk estimates a simulation study: Salt Lake City, UT: University of Utah; 2016. p. 53.
- Egginton JM, Bowles KR, Moyes K, Manley S, Esterling L, Sizemore S, et al. A comprehensive laboratory-based program for classification of variants of uncertain significance in hereditary cancer genes. *Clin Genet* 2014;86:229–37.
- Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. *J Am Statist Assoc* 1958;53:457–81.
- Satagopan JM, Offit K, Foulkes W, Robson ME, Wacholder S, Eng CM, et al. The lifetime risks of breast cancer in Ashkenazi Jewish carriers of BRCA1 and BRCA2 mutations. *Cancer Epidemiol Biomarkers Prev* 2001;10:467–73.
- Surveillance, Epidemiology, and End Results (SEER) Program Populations (1969–2014). National Cancer Institute, DCCPS Surveillance

Hall et al.

- Research Program, Surveillance Systems Branch; 2016. Available from: <https://seer.cancer.gov/>.
33. Rainville I, Hatcher S, Rosenthal E, Larson K, Bernhisel R, Meek S, et al. High risk of breast cancer in women with biallelic pathogenic variants in CHEK2. *Breast Cancer Res Treat* 2020;180:503–9.
 34. Weston VJ, Oldreive CE, Skowronska A, Oscier DG, Pratt G, Dyer MJS, et al. The PARP inhibitor olaparib induces significant killing of ATM-deficient lymphoid tumor cells *in vitro* and *in vivo*. *Blood* 2010; 116:4578–87.
 35. Mateo J, Carreira S, Sandhu S, Miranda S, Mossop H, Perez-Lopez R, et al. DNA-repair defects and olaparib in metastatic prostate cancer. *N Engl J Med* 2015;373:1697–708.
 36. Gatti RA, Tward A, Concannon P. Cancer risk in ATM heterozygotes: a model of phenotypic and mechanistic differences between missense and truncating mutations. *Mol Genet Metab* 1999;68:419–23.
 37. Waddell N, Jonnalagadda J, Marsh A, Grist S, Jenkins M, Hobson K, et al. Characterization of the breast cancer associated ATM 7271T>G (V2424G) mutation by gene expression profiling. *Genes Chromosomes Cancer* 2006;45:1169–81.
 38. Marabelli M, Cheng S-C, Parmigiani G. Penetrance of ATM gene mutations in breast cancer: a meta-analysis of different measures of risk. *Genet Epidemiol* 2016;40:425–31.
 39. Canto MI, Harinck F, Hruban RH, Offerhaus GJ, Poley J-W, Kamel I, et al. International Cancer of the Pancreas Screening (CAPS) Consortium summit on the management of patients with increased risk for familial pancreatic cancer. *Gut* 2013;62: 339–47.
 40. Chaffee KG, Oberg AL, McWilliams RR, Majithia N, Allen BA, Kidd J, et al. Prevalence of germ-line mutations in cancer genes among pancreatic cancer patients with a positive family history. *Genet Med* 2018;20:119–27.
 41. Na R, Zheng SL, Han M, Yu H, Jiang D, Shah S, et al. Germline mutations in ATM and BRCA1/2 distinguish risk for lethal and indolent prostate cancer and are associated with early age at death. *Eur Urol* 2017;71:740–7.
 42. Provenzale D, Gupta S, Ahnen DJ, Blanco AM, Bray TH, Chung DC, et al. NCCN Clinical Practice Guidelines in Oncology, genetic/familial high-risk assessment: colorectal (version 1.2018). In: NCCN Clinical Practice Guidelines in Oncology. Available from: https://www.nccn.org/professionals/physician_gls/pdf/genetics_colon.pdf.
 43. LaDuca H, Polley EC, Yussuf A, Hoang L, Gutierrez S, Hart SN, et al. A clinical guide to hereditary cancer panel testing: evaluation of gene-specific cancer associations and sensitivity of genetic testing criteria in a cohort of 165,000 high-risk patients. *Genet Med* 2020; 22:407–15.
 44. Schlesselman JJ. Case control studies: design, conduct, analysis. New York, NY: Oxford University Press; 1982.

Cancer Prevention Research

Germline Pathogenic Variants in the Ataxia Telangiectasia Mutated (*ATM*) Gene are Associated with High and Moderate Risks for Multiple Cancers

Michael J. Hall, Ryan Bernhisel, Elisha Hughes, et al.

Cancer Prev Res 2021;14:433-440. Published OnlineFirst January 28, 2021.

Updated version	Access the most recent version of this article at: doi: 10.1158/1940-6207.CAPR-20-0448
Supplementary Material	Access the most recent supplemental material at: http://cancerpreventionresearch.aacrjournals.org/content/suppl/2021/01/30/1940-6207.CAPR-20-0448.DC1

Cited articles	This article cites 38 articles, 5 of which you can access for free at: http://cancerpreventionresearch.aacrjournals.org/content/14/4/433.full#ref-list-1
-----------------------	---

E-mail alerts	Sign up to receive free email-alerts related to this article or journal.
----------------------	--

Reprints and Subscriptions	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org .
-----------------------------------	--

Permissions	To request permission to re-use all or part of this article, use this link http://cancerpreventionresearch.aacrjournals.org/content/14/4/433 . Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.
--------------------	--