

Research Article

Black Raspberries Inhibit Intestinal Tumorigenesis in *Apc1638+/-* and *Muc2-/-* Mouse Models of Colorectal CancerXiuli Bi¹, Wenfeng Fang^{1,2}, Li-Shu Wang³, Gary D. Stoner³, and Wancai Yang¹**Abstract**

Freeze-dried black raspberries (BRB) produce chemopreventive effects in a rat model of colon carcinogenesis; however, the mechanisms of inhibition were not determined. Herein, we used two mouse models of human colorectal cancer to determine if dietary BRBs would inhibit colorectal tumor development and to investigate the underlying mechanisms. We found that a 12-week feeding of BRBs significantly inhibited intestinal tumor formation in both models; reducing tumor incidence by 45% and tumor multiplicity by 60% in *Apc1638+/-* mice and tumor incidence and multiplicity by 50% in *Muc2-/-* mice. Mechanistic studies revealed that BRBs inhibit tumor development in *Apc1638+/-* mice by suppressing β -catenin signaling and in *Muc2-/-* mice by reducing chronic inflammation. Intestinal cell proliferation was inhibited by BRBs in both animal models; however, the extent of mucus cell differentiation was not changed in either model. Collectively, our data suggest that BRBs are highly effective in preventing intestinal tumor development in both *Apc1638+/-* and *Muc2-/-* mice through targeting multiple signaling pathways. *Cancer Prev Res*; 3(11); 1443–50. ©2010 AACR.

Introduction

Colorectal cancer is the second leading cause of cancer-related death in men and the third leading cause of cancer death in women in the United States (1). It is also the second most prevalent cancer worldwide. Risk factors for developing colorectal cancer include hereditary predisposition to either familial adenomatous polyposis or nonpolyposis colon cancer, obesity, physical inactivity, smoking, alcohol consumption, and an inadequate intake of vegetables and fruit (2, 3). An important strategy for the prevention of colorectal cancer is endoscopic screening to detect and remove precursor lesions (adenomatous polyps) and to detect early-stage carcinomas. Unfortunately, screening compliance is unacceptably low. Moreover, the currently used chemotherapies for metastatic disease are largely ineffective due to dose-limiting toxicity and acquired chemoresistance; thus, survival rates are poor for patients with metastatic disease.

Chemoprevention refers to the administration of synthetic or naturally occurring agents to block, reverse, or

delay the process of carcinogenesis. For a variety of reasons, including human acceptance and low toxicity, naturally occurring, diet-based agents are ideal for chemopreventive interventions. Berry fruits are widely consumed in our diet and have attracted much attention due to their potential human health benefits. Multiple studies have shown that black raspberries (BRB; *Rubus occidentalis*) exhibit diverse biological properties, including antioxidant, anticancer, antineurodegenerative, and anti-inflammatory activities (4–6). To determine whether BRBs could be useful for the prevention of colorectal cancer, we administered a Western-style diet containing 10% freeze-dried BRBs to two mouse models of human colorectal cancer (i.e., *Apc1638+/-* and *Muc2-/-* mice) for 12 weeks. The *Apc* mouse has one functional allele of the *Apc* gene that, when inactivated, leads to inappropriate signaling of the Wnt/ β -catenin pathway and the spontaneous development of intestinal adenomas (7). *Muc2-/-* mice develop intestinal adenomas and adenocarcinomas in response to chronic inflammation (8, 9). We found that BRBs inhibit intestinal tumor formation in both animal models, and that tumor inhibition was associated with a decrease in β -catenin-induced signaling in *Apc* mice and the inhibition of chronic inflammation in *Muc2-/-* mice.

Materials and Methods**Animals and diets**

The development of the *Apc1638+/-* and *Muc2-/-* mouse models and the methods for genotyping these mice have been described (7–10). After weaning (~3–4 weeks),

Authors' Affiliations: ¹Department of Pathology, University of Illinois at Chicago, Chicago, Illinois; ²The Open Laboratory for Overseas Scientists and Department of Oncology, College of Medicine, Wuhan University, Wuhan, China; and ³Department of Internal Medicine and Comprehensive Cancer Center, The Ohio State University College of Medicine, Columbus, Ohio

Corresponding Author: Wancai Yang, Department of Pathology, University of Illinois at Chicago, Room 130 CSN, 840 South Wood Street, Chicago, IL 60612. Phone: 312-355-4154; Fax: 312-996-7586; E-mail: wyang06@uic.edu.

doi: 10.1158/1940-6207.CAPR-10-0124

©2010 American Association for Cancer Research.

littermates of both mouse strains were randomized to dietary groups and fed either a Western-style (control) diet (11, 12) or the Western-style diet supplemented with 10% (w/w) freeze-dried BRB powder (berry diet). The berry powder was mixed into the Western-style diet by using a Hobart mixer as described before (13). The Western-style diet was formulated on the basis of nutrient density to mimic major risk factors for colon cancer (high in fat and phosphate, and low in calcium and vitamin D). The cornstarch in the berry diet was reduced by 10% to maintain an isocaloric diet. Both the control and berry diets were stored at -20°C before use in the experiment. All animals were housed in plastic cages with filter tops; five mice per cage. The animal room was controlled at $23 \pm 1^{\circ}\text{C}$, $50 \pm 10\%$ humidity, and a 12-hour light/dark cycle. Animals had free access to food and water at all times. Food cups were replenished with fresh diets twice weekly. The animals were housed and maintained in accordance with the recommendations of the University of Illinois at Chicago Animal Use Committee and the American Association of Laboratory Animal Care.

The Western-style diet was purchased from Research Diets, Inc., and the freeze-dried BRB powder was supplied by G. Stoner (Ohio State University Comprehensive Cancer Center). The protocol for procurement, freeze-drying, chemical and microbial characterization, and storage of BRB powder has been described in detail (14).

Histopathology

After 12 weeks of consuming either the control or berry diet, all animals of both mouse strains were sacrificed by CO_2 inhalation followed by cervical dislocation. The entire intestinal tract was removed and opened longitudinally. The contents of the intestine were removed by washing with cold PBS. The full length of the intestinal tract was immediately examined for neoplastic lesions under a dissecting microscope. Tumor location (small intestine or large intestine), incidence (percentage of mice with tumor), multiplicity (number of tumors per mouse), and size (tumor volume) were recorded. Tumors were fixed in 10% buffered formalin. Two fragments, each 0.8 to 1.0 cm in length, of normal-appearing tissue from both the duodenum and colon were placed separately into 10% buffered formalin or snap frozen in liquid nitrogen. Formalin-fixed paraffin-embedded tissues were used for histopathologic (hematoxylin and eosin staining) and immunohistochemical staining. Frozen tissues were used to prepare frozen sections for histopathology or for biochemical studies.

Proliferative cell nuclear antigen (PCNA) staining was used to evaluate cell proliferation. PCNA-stained and unstained epithelial cells from about 25 well-oriented crypts per intestine per mouse in both diet groups were counted, and the percentage of PCNA-positive cells was calculated. Alcian blue staining was performed to evaluate goblet cell differentiation. The percentage of Alcian blue-positive cells in the intestinal crypts of each mouse was determined in the same manner as for PCNA-positive cells. All procedures, including evaluation of tumorigenesis, PCNA, and

Alcian blue staining and scoring, have been standardized in our laboratory as described previously (10, 15–19).

Intestinal epithelial cell isolation and quantitative real-time reverse transcriptase-PCR

Intestinal epithelial cells from both mouse strains were isolated by incubating opened mouse intestine from colon or from the combination of duodenum and jejunum, respectively, in 15 mmol/L EDTA buffer at 37°C for 30 minutes as described previously (15). Total RNA was extracted from epithelial cell pellets obtained from three individual mice per strain per diet group. The quality and quantity of RNA was determined using a NanoDrop Spectrophotometer (Thermo Scientific). Quantitative real-time PCR analysis was done using the ABI Prism 7900-HT sequence detection system (96 wells; Applied Biosystems) as described (15). The following primers were designed for mouse cytokine analysis and were synthesized by Sigma Oligo: cyclooxygenase-2 (COX-2), forward 5'-TGAGCAACTATTCCTCAA CCAGC-3' and reverse 5'-GCACGTAGTCTTCGATCACTATC-3'; tumor necrosis factor- α (TNF- α), forward 5'-CCCTCACACTCAGATCATCTTCT-3' and reverse 5'-GCTACGACGTGGCTACAG-3'; interleukin (IL)-1, forward 5'-GCAACTGTTCTGAACTCAACT-3' and reverse 5'-ATCTTTTGGGGTCCGTCAACT-3'; IL-6, forward 5'-TAGTCCTTCCTACCCCAATTTCC-3' and reverse 5'-TTGGTCCTTAGCCACTCCTC-3'; IL-10, forward 5'-GCTCTTACTGACTGGCATGAG-3' and reverse 5'-CCGAGCTCTAGGAGCATGTG-3'. β -Actin was used as internal control (15).

Western blotting analyses and immunohistochemical staining

In addition to RNA, protein was extracted from each of the above-described intestinal epithelial cell pellets. In brief, the pellets were washed twice with ice-cold PBS and incubated on ice for 15 minutes in cell lysis buffer (Cell Signaling). After brief sonication, the cell lysate was centrifuged at 13,500 rpm for 15 minutes at 4°C . The supernatant was fractionated by electrophoresis in a 12% SDS-polyacrylamide gel and transferred to a nitrocellulose membrane. The following primary antibodies were used for immunoblotting: anti-p21 (1:500), anti-p27 (1:1,000), anti- β -catenin (1:1,000), anti-E-cadherin (1:1,000), anti-c-myc (1:1,000), anti-cdk4 (1:1,000), anti-cyclin D1 (1:1,000; Santa Cruz Biotechnology), and anti- β -actin (1:10,000; Sigma). Signals were detected by an enhanced chemiluminescence technique (Amersham Life Science).

Protein expression in the intestine was also analyzed by immunohistochemistry, as we described previously (15). Briefly, formalin-fixed and paraffin-embedded mouse intestinal tissues were sectioned, deparaffinized, and dehydrated. Five animals were analyzed from each of the control and berry diet groups. To block endogenous peroxidase, the sections were incubated in H_2O_2 (0.3%)/methanol for 20 minutes. Sections were then incubated with 10% normal goat serum to block nonspecific antibody binding. To expose epitopes for immunodetection,

sections were treated in a steamer for 20 minutes in citrate buffer (pH 6.1). The procedure for PCNA staining was as described by the manufacturer, and the percentage of PCNA-stained cells was determined as described above.

Statistical analysis

The χ^2 test was used for tumor incidence analysis, and the Student's *t* test was used for tumor multiplicity and other quantification [including PCNA scoring and quantitative reverse transcriptase-PCR (RT-PCR)] analysis. *P* < 0.05 indicated a significant difference.

Results

Dietary BRBs inhibit intestinal tumorigenesis in *Apc1638+/-* mice

After 12 weeks of treatment with the control and BRB diets, all *Apc1638+/-* mice were sacrificed and their intestinal tumors were quantitated. All (11 of 11) of the *Apc1638+/-* mice fed the Western-style control diet developed small intestinal tumors, with an average of 2.33 tumors per mouse (Fig. 1). However, only 55% (6 of 11) of mice fed the 10% BRB-supplemented diet developed intestinal tumors, and the tumor multiplicity was reduced significantly (*P* < 0.01) to 0.91 tumors per mouse (Fig. 1). The tumor sizes were slightly reduced by BRBs in the *Apc1638+/-* mice, but the reduction was not significant (data not shown), which could be due to short-term (12 weeks) feeding.

Dietary BRBs inhibit tumor formation in *Muc2-/-* mice

Unlike *Apc1638+/-* mice, *Muc2-/-* mice developed tumors throughout the entire intestine, principally in the colon and rectum. As shown in Fig. 2, in *Muc2-/-* mice fed the control diet, 44% (8 of 18) developed small intestinal tumors and 100% (18 of 18) developed colon and rectal tumors. However, feeding the BRB diet reduced the incidence of small intestinal tumors by ~30% [44% in the Western-style diet versus 14% (3 of 21) in the BRB diet] and the multiplicity from 0.67 to 0.14 tumors per mouse (*P* < 0.05). Similarly, in the colon, tumor incidence was decreased by 24% (100% versus 76%; 16 of 21) and the multiplicity was reduced remarkably from 3.83 to 1.60 tumors per mouse (*P* < 0.001). In the rectum, tumor multiplicity was reduced dramatically from 1.0 to 0.37 per mouse (*P* < 0.001) and tumor incidence was decreased by ~62%, from 100% (18 of 18) to 38% (8 of 21). The difference of tumor sizes in the control and BRBs groups was not significant in the *Muc2-/-* mice (data not shown).

Tumor inhibition by BRBs is linked to reduced intestinal cell proliferation and not differentiation

We investigated the effects of BRBs on intestinal cell proliferation in both mouse strains. The duodenum, the principal site of tumor formation in *Apc1638+/-* mice, and the colon, the major site of tumor formation in *Muc2-/-* mice, were stained for PCNA. BRBs significantly inhibited intestinal cell proliferation in the duodenum of




Fig. 1. BRBs significantly inhibit intestinal tumor incidence (A) and multiplicity (B; mean \pm SD) in *Apc1638+/-* mice at 12 weeks. (*, *P* < 0.05; **, *P* < 0.01, compared with mice fed a Western-style diet; 11 mice per group were studied).

Apc1638+/- mice (Fig. 3A) and in the colon of *Muc2-/-* mice (Fig. 3B). However, the berry diet did not alter intestinal cell differentiation as measured by Alcian blue staining for goblet cell identification (data not shown).

BRB-induced tumor inhibition in *Apc1638+/-* mice is associated with reduced β -catenin signaling and a modest suppression of cell cytokines

Aberrant β -catenin expression is a common feature of APC mutation-induced tumorigenesis in the small intestine of *Apc+/-* mice and in human colorectal cancer (20–22). We determined, therefore, the expression levels of β -catenin and its downstream targets in normal intestinal mucosa of *Apc1638+/-* mice fed either the control or 10% BRB diet. Proteins extracted from mouse intestinal

epithelial cells were evaluated by Western blotting. The level of β -catenin was dramatically decreased, and c-Myc and cyclin D1, both downstream of β -catenin, were modestly decreased in BRB-fed mice. In contrast, the cyclin-dependent kinase inhibitor, p27kip1, was increased in BRB-fed mice when compared with mice fed the control diet (Fig. 4A).

To determine whether BRBs produced anti-inflammatory effects in *Apc*^{1638+/-} mice, we used quantitative RT-PCR to evaluate changes in a set of inflammatory biomarkers in intestinal epithelial cells. The mRNA levels of COX-2 and of the proinflammatory cytokines, TNF- α , IL-6, and IL-10, were modestly decreased, whereas IL-1 was slightly increased (Fig. 4B). None of these changes were significant.

BRB-induced tumor inhibition in *Muc2*^{-/-} mice is associated with inhibition of chronic inflammation but not with β -catenin signaling

We have reported that tumor formation in *Muc2*^{-/-} mice is associated with chronic inflammation; β -catenin is not involved (8, 9). To evaluate the effects of BRBs




Fig. 2. BRBs significantly inhibit intestinal tumor incidence (A) and multiplicity (B; mean \pm SD) in *Muc2*^{-/-} mice at 12 weeks. (*, $P < 0.05$; ***, $P < 0.001$, compared with mice fed a Western-style diet. There were 18 mice in the Western-style diet group and 21 mice in the BRBs groups, respectively).




Fig. 3. BRBs inhibit intestinal cell proliferation (assayed by PCNA staining) in the *Apc*^{1638+/-} mouse small intestine (A) and the *Muc2*^{-/-} mouse colon (B). (*, $P < 0.05$, compared with mice fed a Western-style diet).

on inflammation in *Muc2*^{-/-} mice, we determined the mRNA expression levels of cytokines in the supernatants of whole colonic epithelial cells. COX-2 and all proinflammatory cytokines (i.e., TNF- α , IL-1, IL-6, and IL-10) were significantly decreased by BRBs ($P < 0.05$ or $P < 0.01$; Fig. 5A).


We also investigated changes in proteins involved in β -catenin signaling in both the small intestine and colon of *Muc2*^{-/-} mice. Unlike in *Apc*^{1638+/-} mice, the β -catenin signaling pathway in the small intestine of *Muc2*^{-/-} mice was not influenced by berry treatment (Fig. 5B). However, E-cadherin and another cyclin-dependent kinase inhibitor, p21^{WAF1/Cip1}, were induced by BRBs in the colon of *Muc2*^{-/-} mice (Fig. 5C).

Discussion

BRBs are known to contain multiple compounds with chemopreventive potential (4, 5, 13, 14). Previous studies have shown the protective effects of dietary freeze-dried BRBs on the occurrence of chemically induced tumors in rodents, including tumors of the colon (4, 14, 23). The molecular events involved in the inhibition by BRBs of chemically induced tumorigenesis in the rat esophagus have been investigated in detail (4, 5, 24–28); however, there is very little mechanistic information at the molec-

ular level for BRBs in the rodent colon (23). Using two distinctive mouse models, *Apc1638+/-* and *Muc2-/-* mice, each of which has unique molecular events for study, we found that dietary BRBs were effective in inhibiting intestinal tumorigenesis. In *Apc1638+/-* mice, this inhibition likely occurred through the effects of berries on aberrant β -catenin signaling, whereas in *Muc2-/-* mice, it was linked to inhibition of chronic inflammation and increase in the expression of E-cadherin and p21. Collectively, these data suggest that BRBs produce a broad range of protective effects in the intestine, colon, and rectum.

Fig. 4. Alterations in β -catenin signaling (A) and inflammatory factors (B; mean \pm SD) in *Apc+/-* mice by BRBs as assayed by Western blotting and real-time RT-PCR, respectively. Protein and RNA were extracted from intestinal epithelial cells from three individual mice per group.






Fig. 5. Alterations in inflammatory factors (A; mean \pm SD) and molecular changes (B and C) by BRBs in *Muc2*^{-/-} mice analyzed by real-time RT-PCR and Western blotting, respectively. RNA was extracted from colon tissues, and the protein for Western blotting was extracted from small intestinal epithelial cells (B) and colonic epithelial cells (C) from three individual mice per group. (*, $P < 0.05$; **, $P < 0.01$, compared with the mice fed a Western-style diet).

This is in agreement with our previous investigations in which dietary BRBs were shown to produce a genome-wide protective effect on the expression of genes associated with the development of tumors in carcinogen-treated rat esophagus (24, 28).

Mutations in the *Apc* gene are frequently observed in human colorectal cancers, and β -catenin activation is an early event of *APC* mutation-initiated colorectal carcinogenesis. *Apc* mutations in the mouse lead to the accumulation of cytoplasmic β -catenin, its subsequent nuclear translocation, and the resultant activation of aberrant TCF4-*c-Myc* signaling. Thus, Wnt/ β -catenin signaling is

a promising target for chemoprevention and chemotherapy of colorectal cancer (29, 30). Our previous studies have shown that sulindac, a nonsteroidal anti-inflammatory drug, and the naturally occurring compound 20(S)-25-OCH₃-PPD, derived from the leaves of *Panax notoginseng*, both inhibit human colorectal cancer cell proliferation and promote apoptosis by targeting the Wnt/ β -catenin signaling pathway (31, 32). Here, we show that BRBs also target β -catenin signaling in *Apc1638*^{+/-} mice; intriguingly, as shown in Fig. 4A, β -catenin protein levels in nontumor epithelial cells were upregulated by the Western high-risk diet (9, 33) and were driven back to normal levels by

BRBs. Our data suggest that suppression of the Wnt signaling pathway is another mechanism by which BRBs inhibit tumorigenesis.

Muc2^{-/-} mice spontaneously develop intestinal adenomas that frequently progress to invasive adenocarcinomas when the animals reach 6 months to 1 year in age. Tumor development in the intestine of *Muc2*^{-/-} mice is associated with activation of inflammation-related pathways (8, 9). Recent studies in our laboratory indicate that the Western-style diet markedly accelerates colorectal cancer formation in *Muc2*^{-/-} mice; that is, tumors arise within 3 months.⁴ Herein, we found that BRBs significantly inhibit colorectal tumor formation in *Muc2*^{-/-} mice fed the Western-style diet. The berries dramatically decreased mRNA expression levels of the proinflammatory cytokines, IL-1, IL-6, IL-10, TNF- α , and COX-2. Interestingly, E-cadherin and p21 were significantly increased in the colon, and β -catenin was not changed. The differences of BRB inhibition of β -catenin in the two mouse models could be caused by the differential interaction between BRBs and the *Apc* or *Muc2* genes. Whether there is a gene-diet interaction or how this interaction affects the efficacy of BRB tumor inhibition is under investigation.

E-cadherin, a protein that mediates cell-cell adhesion through calcium-dependent homophilic interactions in the extracellular domain, plays an important role in epithelial-mesenchymal transition. p21 has been shown to regulate E-cadherin (34), but the mechanism(s) of its regulation are not clear. It is well known that E-cadherin binds to β -catenin, and the E-cadherin/ β -catenin complex plays an important role in cell proliferation and differentiation during development (35). The observation that E-cadherin, but not β -catenin, was increased in the intestine of berry-fed animals in the present study could have been due to the formation of a complex between β -catenin and E-cadherin. Such a complex could limit the translocation of β -catenin from the cell membrane into the nucleus, resulting in the modest effects

we observed of the berry diet on c-Myc and cyclin D1 expression levels in the intestine.

Epithelial-mesenchymal transition occurs at the invasive front of tumors, the same site where tumors are infiltrated by tumor-associated macrophages. The decreased expression of macrophage-associated TNF- α in berry-fed animals could result in the stabilization of Snail, a transcription factor that represses the expression of E-cadherin (36, 37). Currently, we are investigating whether the enhanced E-cadherin expression in berry-fed animals is the result of a decreased stability of Snail mediated by TNF- α .

In summary, the present study shows that dietary BRBs are effective in inhibiting intestinal tumor formation in the *Apc*¹⁶³⁸/⁺/⁻ and *Muc2*^{-/-} mouse models of intestinal and colorectal cancer. Tumor inhibition is associated with reduced cell proliferation, suppression of the β -catenin signaling and chronic inflammation pathways, and increase in cellular E-cadherin. Our data add importantly to the existing knowledge on the molecular mechanisms of BRB-mediated chemoprevention and provide additional rationale for assessing the potential chemopreventive effects of BRBs for human colorectal cancer.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

We thank Dr. Leonard Augenlicht (Albert Einstein Cancer Center, Bronx, NY) for providing the *Apc*¹⁶³⁸/⁺/⁻ mice, Dr. Anna Velcich (Albert Einstein Cancer Center, Bronx, NY) for providing the *Muc2*^{-/-} mice, and the University of Illinois at Chicago Cancer Center Histopathology Core Facility for technical assistance.

Grant Support

Faculty Startup Fund (W. Yang) from the Department of Pathology, University of Illinois at Chicago, and National Cancer Institute grants CA112081 (W. Yang) and CA103180 (G. Stoner).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received 06/01/2010; revised 07/31/2010; accepted 09/06/2010; published OnlineFirst 11/02/2010.

⁴ Fang and Yang et al., unpublished data.

References

- Kelloff GJ, Lippman SM, Dannenberg AJ, et al. Progress in chemoprevention drug development: the promise of molecular biomarkers for prevention of intraepithelial neoplasia and cancer—a plan to move forward. *Clin Cancer Res* 2006;12:3661–97.
- Norat T, Bingham S, Ferrari P, et al. Meat, fish, and colorectal cancer risk: the European Prospective Investigation into Cancer and Nutrition. *J Natl Cancer Inst* 2005;97:906–16.
- Riboli E, Norat T. Epidemiologic evidence of the protective effect of fruit and vegetables on cancer risk. *Am J Clin Nutr* 2003;78:559–69S.
- Stoner GD. Foodstuffs for preventing cancer: the preclinical and clinical development of berries. *Cancer Prev Res (Phila)* 2009;2:187–94.
- Wang LS, Hecht SS, Carmella SG, et al. Anthocyanins in black raspberries prevent esophageal tumors in rats. *Cancer Prev Res (Phila)* 2009;2:84–93.
- Seeram NP, Adams LS, Zhang Y, et al. Blackberry, black raspberry, blueberry, cranberry, red raspberry, and strawberry extracts inhibit growth and stimulate apoptosis of human cancer cells *in vitro*. *J Agric Food Chem* 2006;54:9329–39.
- Fodde R, Edelmann W, Yang K, et al. A targeted chain-termination mutation in the mouse *Apc* gene results in multiple intestinal tumors. *Proc Natl Acad Sci U S A* 1994;91:8969–73.
- Velcich A, Yang W, Heyer J, et al. Colorectal cancer in mice genetically deficient in the mucin *Muc2*. *Science (New York, NY)* 2002;295:1726–9.
- Yang K, Popova NV, Yang WC, et al. Interaction of *Muc2* and *Apc* on Wnt signaling and in intestinal tumorigenesis: potential role of chronic inflammation. *Cancer Res* 2008;68:7313–22.
- Yang W, Velcich A, Lozonoschi I, et al. Inactivation of p21WAF1/cip1

- enhances intestinal tumor formation in *Muc2*^{-/-} mice. *Am J Pathol* 2005;166:1239–46.
11. Newmark HL, Lipkin M, Maheshwari N. Colonic hyperplasia and hyperproliferation induced by a nutritional stress diet with four components of Western-style diet. *J Natl Cancer Inst* 1990;82:491–6.
 12. Newmark HL, Lipkin M, Maheshwari N. Colonic hyperproliferation induced in rats and mice by nutritional-stress diets containing four components of a human Western-style diet (series 2). *Am J Clin Nutr* 1991;54:209–14S.
 13. Kresty LA, Morse MA, Morgan C, et al. Chemoprevention of esophageal tumorigenesis by dietary administration of lyophilized black raspberries. *Cancer Res* 2001;61:6112–9.
 14. Stoner GD, Wang LS, Casto BC. Laboratory and clinical studies of cancer chemoprevention by antioxidants in berries. *Carcinogenesis* 2008;29:1665–74.
 15. Bi X, Tong C, Dockendorff A, et al. Genetic deficiency of decorin causes intestinal tumor formation through disruption of intestinal cell maturation. *Carcinogenesis* 2008;29:1435–40.
 16. Tong C, Yin Z, Song Z, et al. c-Jun NH₂-terminal kinase 1 plays a critical role in intestinal homeostasis and tumor suppression. *Am J Pathol* 2007;171:297–303.
 17. Yang W, Bancroft L, Liang J, Zhuang M, Augenlicht LH. p27kip1 in intestinal tumorigenesis and chemoprevention in the mouse. *Cancer Res* 2005;65:9363–8.
 18. Yang W, Bancroft L, Nicholas C, Lozonschi I, Augenlicht LH. Targeted inactivation of p27kip1 is sufficient for large and small intestinal tumorigenesis in the mouse, which can be augmented by a Western-style high-risk diet. *Cancer Res* 2003;63:4990–6.
 19. Yang W, Mathew J, Velcich A, et al. Targeted inactivation of the p21 (*WAF1/cip1*) gene enhances *Apc*-initiated tumor formation and the tumor-promoting activity of a Western-style high-risk diet by altering cell maturation in the intestinal mucosal. *Cancer Res* 2001;61:565–9.
 20. Korinek V, Barker N, Morin PJ, et al. Constitutive transcriptional activation by a β -catenin-Tcf complex in *APC*^{-/-} colon carcinoma. *Science* 1997;275:1784–7.
 21. Morin PJ, Sparks AB, Korinek V, et al. Activation of β -catenin-Tcf signaling in colon cancer by mutations in β -catenin or *APC*. *Science* 1997;275:1787–90.
 22. Rubinfeld B, Albert I, Porfiri E, Munemitsu S, Polakis P. Loss of β -catenin regulation by the *APC* tumor suppressor protein correlates with loss of structure due to common somatic mutations of the gene. *Cancer Res* 1997;57:4624–30.
 23. Harris GK, Gupta A, Nines RG, et al. Effects of lyophilized black raspberries on azoxymethane-induced colon cancer and 8-hydroxy-2'-deoxyguanosine levels in the Fischer 344 rat. *Nutr Cancer* 2001;40:125–33.
 24. Wang L-S, Dombkowski AA, Rocha C, et al. Effects of black raspberries on late events in *N*-nitrosomethylbenzylamine-induced rat esophageal carcinogenesis as determined by DNA microarray. *Mol Carcinogen*. In press 2010.
 25. Stoner GD, Chen T, Kresty LA, Aziz RM, Reinemann T, Nines R. Protection against esophageal cancer in rodents with lyophilized berries: potential mechanisms. *Nutr Cancer* 2006;54:33–46.
 26. Chen T, Hwang H, Rose ME, Nines RG, Stoner GD. Chemopreventive properties of black raspberries in *N*-nitrosomethylbenzylamine-induced rat esophageal tumorigenesis: down-regulation of cyclooxygenase-2, inducible nitric oxide synthase, and c-Jun. *Cancer Res* 2006;66:2853–9.
 27. Chen T, Rose ME, Hwang H, Nines RG, Stoner GD. Black raspberries inhibit *N*-nitrosomethylbenzylamine (NMBM)-induced angiogenesis in rat esophagus parallel to the suppression of COX-2 and iNOS. *Carcinogenesis* 2006;27:2301–7.
 28. Stoner GD, Dombkowski AA, Reen RK, et al. Carcinogen-altered genes in rat esophagus positively modulated to normal levels of expression by both black raspberries and phenylethyl isothiocyanate. *Cancer Res* 2008;68:6460–7.
 29. Herbst A, Kolligs FT. Wnt signaling as a therapeutic target for cancer. *Methods Mol Biol (Clifton, NJ)* 2007;361:63–91.
 30. Luo J, Chen J, Deng ZL, et al. Wnt signaling and human diseases: what are the therapeutic implications? *Lab Invest J Tech Methods Pathol* 2007;87:97–103.
 31. Bi X, Zhao Y, Fang W, Yang W. Anti-cancer bioactivities of *Panax notoginseng* extract 20(S)-25-OCH₃-PPD: targeting β -catenin signaling. *Clin Exp Pharmacol Physiol* 2009;36:1074–8.
 32. Han A, Song Z, Tong C, et al. Sulindac suppresses β -catenin expression in human cancer cells. *Eur J Pharmacol* 2008;583:26–31.
 33. Yang K, Kurihara N, Fan K, et al. Dietary induction of colonic tumors in a mouse model of sporadic colon cancer. *Cancer Res* 2008;68:7803–10.
 34. Mueller S, Cadenas E, Schonthal AH. p21WAF1 regulates anchorage-independent growth of HCT116 colon carcinoma cells via E-cadherin expression. *Cancer Res* 2000;60:156–63.
 35. Wheelock MJ, Johnson KR. Cadherins as modulators of cellular phenotype. *Ann Rev Cell Dev Biol* 2003;19:207–35.
 36. Yang CC, Wolf DA. Inflamed snail speeds metastasis. *Cancer Cell* 2009;15:355–7.
 37. Wu Y, Deng J, Rychahou PG, Qiu S, Evers BM, Zhou BP. Stabilization of snail by NF- κ B is required for inflammation-induced cell migration and invasion. *Cancer Cell* 2009;15:416–28.

Cancer Prevention Research

Black Raspberries Inhibit Intestinal Tumorigenesis in *Apc* 1638+/- and *Muc2*-/- Mouse Models of Colorectal Cancer

Xiuli Bi, Wenfeng Fang, Li-Shu Wang, et al.

Cancer Prev Res 2010;3:1443-1450. Published OnlineFirst November 2, 2010.

Updated version Access the most recent version of this article at:
doi:[10.1158/1940-6207.CAPR-10-0124](https://doi.org/10.1158/1940-6207.CAPR-10-0124)

Cited articles This article cites 36 articles, 16 of which you can access for free at:
<http://cancerpreventionresearch.aacrjournals.org/content/3/11/1443.full#ref-list-1>

Citing articles This article has been cited by 5 HighWire-hosted articles. Access the articles at:
<http://cancerpreventionresearch.aacrjournals.org/content/3/11/1443.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link <http://cancerpreventionresearch.aacrjournals.org/content/3/11/1443>. Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.