

21. Delker DA, McGettigan BM, Kanth P, Pop S, Neklason DW, Bronner MP, et al. RNA sequencing of sessile serrated colon polyps identifies differentially expressed genes and immunohistochemical markers. *PLoS One* 2014;9:e88367.
22. Lieberman DA, Rex DK, Winawer SJ, Giardiello FM, Johnson DA, Levin TR, et al. United States Multi-Society Task Force on Colorectal Cancer. Guidelines for colonoscopy surveillance after screening and polypectomy: a consensus update by the US Multi-Society Task Force on Colorectal Cancer. *Gastroenterology* 2012;143:844–57.
23. Folkers ME, Delker DA, Maxwell CI, Nelson CA, Schwartz JJ, Nix DA, et al. ENCODE tiling array analysis identifies differentially expressed annotated and novel 5' capped RNAs in hepatitis C infected liver. *PLoS One* 2011;6:e14697.
24. Papic N, Maxwell CI, Delker DA, Liu S, Heale BS, Hagedorn CH. RNA-sequencing analysis of 5' capped RNAs identifies many new differentially expressed genes in acute hepatitis C virus infection. *Viruses* 2012;4:581–612.
25. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-Seq data with DESeq2. *Genome Biol* 2014;15:550.
26. Comprehensive molecular characterization of human colon and rectal cancer. Cancer Genome Atlas Network. *Nature* 2012;487:330–7.
27. National Institutes of Health. TCGA data portal. Available from: <https://tcga-data.nci.nih.gov/tcga/tcgaHome2.jsp>.
28. Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, et al. Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signali* 2013;6:p11.
29. Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov* 2012;2:401–4.
30. Ciriello G, Cerami E, Sander C, Schultz N. Mutual exclusivity analysis identifies oncogenic network modules. *Genome Res* 2012;22:398–406.
31. Caruso M1, Fung KY2, Moore J3, Brierley GV, Cosgrove LJ, Thomas M, et al. Claudin-1 expression is elevated in colorectal cancer precursor lesions harboring the BRAF V600E. *Transl Oncol* 2014;7:456–63.
32. DiSario JA, Foutch PG, Mai HD, Pardy K, Manne RK. Prevalence and malignant potential of colorectal polyps in asymptomatic, average-risk men. *Am J Gastroenterol* 1995;90:941–5.
33. Weston AP, Campbell DR. Diminutive colonic polyps: histopathology, spatial distribution, concomitant significant lesions, and treatment complications. *Am J Gastroenterol* 1995;90:24–8.
34. Popovici V, Budinska E, Tejpar S, Weinrich S, Estrella H, Hodgson G, et al. Identification of a poor-prognosis BRAF-mutant-like population of patients with colon cancer. *J Clin Oncol* 2012;30:1288–95.
35. Guinney J, Dienstmann R, Wang X, de Reynies A, Schlicker A, Soneson C, et al. The consensus molecular subtypes of colorectal cancer. *Nat Med* 2015;21:1350–6.
36. De Sousa EMelo F, Wang X, Jansen M, Fessler E, Trinh A, de Rooij LP, et al. Poor-prognosis colon cancer is defined by a molecularly distinct subtype and develops from serrated precursor lesions. *Nat Med* 2013;19:614–8.
37. Merzdorf CS. Emerging roles for zic genes in early development. *Dev Dyn* 2007;236:922–40.
38. Sanek NA, Taylor AA, Nyholm MK, Grinblat Y. Zebrafish *zic2a* patterns the forebrain through modulation of Hedgehog-activated gene expression. *Development* 2009;136:3791–800.
39. Marchini S, Poynor E, Barakat RR, Clivio L, Cinquini M, Fruscio R, et al. The zinc finger gene *ZIC2* has features of an oncogene and its overexpression correlates strongly with the clinical course of epithelial ovarian cancer. *Clin Cancer Res* 2012;18:4313–24.
40. Chan DW, Liu VW, Leung LY, Yao KM, Chan KK, Cheung AN, et al. *Zic2* synergistically enhances Hedgehog signalling through nuclear retention of Gli1 in cervical cancer cells. *J Pathol* 2011;225:525–34.
41. Ma Y, Machesky LM. Fascin1 in carcinomas: Its regulation and prognostic value. *Int J Cancer* 2015;137:2534–44.
42. Conesa-Zamora P, García-Solano J, García-García F, Turpin Mdel C, Trujillo-Santos J, Torres-Moreno D, et al. Expression profiling shows differential molecular pathways and provides potential new diagnostic biomarkers for colorectal serrated adenocarcinoma. *Int J Cancer* 2013;132:297–307.
43. Stahl R, Walcher T, De Juan Romero C, Pilz GA, Cappello S, Imler M, et al. *Tmp1* regulates expansion and folding of the mammalian cerebral cortex by control of radial glial fate. *Cell* 2013;153:535–49.
44. Owens SR, Chiose SI, Kuan SF. Selective expression of gastric mucin *MUC6* in colonic sessile serrated adenoma but not in hyperplastic polyp aids in morphological diagnosis of serrated polyps. *Mod Pathol* 2008;21:660–9.
45. Walsh MD, Clendenning M, Williamson E, Pearson SA, Walters RJ, Nagler B, et al. Expression of *MUC2*, *MUC5AC*, *MUC5B*, and *MUC6* mucins in colorectal cancers and their association with the CpG island methylator phenotype. *Mod Pathol* 2013;26:1642–56.
46. Neuhaus J, Schiffer E, von Wilcke P, Bauer HW, Leung H, Siwy J, et al. Seminal plasma as a source of prostate cancer peptide biomarker candidates for detection of indolent and advanced disease. *PLoS One* 2013;8:e67514.
47. Abe M, Watanabe N, McDonell N, Takato T, Ohira M, Nakagawara A, et al. Identification of genes targeted by CpG island methylator phenotype in neuroblastomas, and their possible integrative involvement in poor prognosis. *Oncology* 2008;74:50–60.
48. Hawthorn L, Lan L, Mojica W. Evidence for field effect cancerization in colorectal cancer. *Genomics* 2014;103:211–21.
49. Lochhead P, Chan AT, Nishihara R, Fuchs CS, Beck AH, Giovannucci E, et al. Etiologic field effect: reappraisal of the field effect concept in cancer predisposition and progression. *Mod Pathol* 2015;28:14–29.
50. Chen LC, Hao CY, Chiu YS, Wong P, Melnick JS, Brotman M, et al. Alteration of gene expression in normal-appearing colon mucosa of APC(min) mice and human cancer patients. *Cancer Res* 2004;64:3694–700.
51. Huang CS, Farraye FA, Yang S, O'Brien MJ. The clinical significance of serrated polyps. *Am J Gastroenterol* 2011;106:229–40.
52. Donehower LA, Creighton CJ, Schultz N, Shinbrot E, Chang K, Gunaratne PH, et al. MLH1-silenced and non-silenced subgroups of hypermutated colorectal carcinomas have distinct mutational landscapes. *J Pathol* 2013;229:99–110.
53. Burnett-Hartman AN, Newcomb PA, Potter JD, Passarelli MN, Phipps AI, Wurscher MA, et al. Genomic aberrations occurring in subsets of serrated colorectal lesions but not conventional adenomas. *Cancer Res* 2013;73:2863–72.

Cancer Prevention Research

Gene Signature in Sessile Serrated Polyps Identifies Colon Cancer Subtype

Priyanka Kanth, Mary P. Bronner, Kenneth M. Boucher, et al.

Cancer Prev Res 2016;9:456-465. Published OnlineFirst March 29, 2016.

Updated version	Access the most recent version of this article at: doi:10.1158/1940-6207.CAPR-15-0363
Supplementary Material	Access the most recent supplemental material at: http://cancerpreventionresearch.aacrjournals.org/content/suppl/2016/03/29/1940-6207.CAPR-15-0363.DC1

Cited articles	This article cites 50 articles, 11 of which you can access for free at: http://cancerpreventionresearch.aacrjournals.org/content/9/6/456.full#ref-list-1
-----------------------	--

Citing articles	This article has been cited by 4 HighWire-hosted articles. Access the articles at: http://cancerpreventionresearch.aacrjournals.org/content/9/6/456.full#related-urls
------------------------	---

E-mail alerts	Sign up to receive free email-alerts related to this article or journal.
----------------------	--

Reprints and Subscriptions	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org .
-----------------------------------	--

Permissions	To request permission to re-use all or part of this article, use this link http://cancerpreventionresearch.aacrjournals.org/content/9/6/456 . Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.
--------------------	--